

Genetic control of branching patterns in grass inflorescences

Elizabeth A. Kellogg ^{1,*†}

¹ Donald Danforth Plant Science Center, St Louis, Missouri 63132, USA

*Author for correspondence: ekellogg@danforthcenter.org

†Senior author

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plcell>) is: Elizabeth A. Kellogg (ekellogg@danforthcenter.org).

Abstract

Inflorescence branching in the grasses controls the number of florets and hence the number of seeds. Recent data on the underlying genetics come primarily from rice and maize, although new data are accumulating in other systems as well. This review focuses on a window in developmental time from the production of primary branches by the inflorescence meristem through to the production of glumes, which indicate the transition to producing a spikelet. Several major developmental regulatory modules appear to be conserved among most or all grasses. Placement and development of primary branches are controlled by conserved auxin regulatory genes. Subtending bracts are repressed by a network including *TASSELSEATH4*, and axillary branch meristems are regulated largely by signaling centers that are adjacent to but not within the meristems themselves. Gradients of *SQUAMOSA-PROMOTER BINDING*-like and *APETALA2*-like proteins and their microRNA regulators extend along the inflorescence axis and the branches, governing the transition from production of branches to production of spikelets. The relative speed of this transition determines the extent of secondary and higher order branching. This inflorescence regulatory network is modified within individual species, particularly as regards formation of secondary branches. Differences between species are caused both by modifications of gene expression and regulators and by presence or absence of critical genes. The unified networks described here may provide tools for investigating orphan crops and grasses other than the well-studied maize and rice.

Introduction

Grass dominated ecosystems cover ~40% of the Earth's land surface (Lehmann et al., 2019; Griffith et al., 2020) and provide over 50% of the world's calories, whether consumed directly or fed to animals which are then consumed by humans (FAO, 2003). Central to the ecological and economic roles of grasses is the inflorescence, the complex set of flowers that produces seeds. The combined number and size of seeds contribute to higher fitness in the wild and higher yield in cultivation. Accordingly, inflorescence structure and flower/seed production have been the target of both natural and human selection.

Grass inflorescence development is often described by invoking shifting meristem identity. As a shoot apical

meristem (SAM) producing leaves on its flanks changes to producing bracts, branches, and floral meristems (FMs), it is described as acquiring inflorescence meristem (IM) identity. Within the inflorescence, similar shifts specify branch meristem (BM) identity and FM identity. In addition, in grasses a distinct structure, the spikelet (a tiny spike), is interpolated developmentally between the BM and FM and arises from a spikelet meristem (SM). The metaphor of meristem identity implies that the meristem is itself somehow autonomous and distinct from both the surrounding cells and also from other meristems on the plant.

In a thought-provoking paper, Whipple (2017) observed that the concept of meristem identity also implies the existence of selector genes whose presence confers particular

characteristics on a meristem. While such selector genes are known for FMs, they have been elusive for other meristem types. Instead, he notes that the boundary between a meristem and its subtending bracts has emerged as an important signaling center (Whipple, 2017), shifting the metaphor from the meristem as a master controller of its own fate to the meristem as an emergent structure, the result of disparate inputs and outputs.

Another common metaphor is that of a developmental switch, a gene being either on or off. However, gene regulation is often quantitative, more like a rheostat than a binary switch, leading to metaphors of phase change (e.g. Kyoizuka, 2014) and gradual transitions from one state to another. At the beginning and end of the transition, a structure may be recognizable as an IM, BM, or SM, but the boundaries between them may not be sharp, although a gradual transition might ultimately trigger a switch. These metaphors—identity and signaling centers, switches, and rheostats—currently co-exist productively and are themes that occur throughout this review.

This review focuses on a narrow but critical developmental window, from the production of primary BMs by the IM through to specification of spikelets. These are processes that vary extensively in response to natural (evolutionary) and human (agricultural) selection. Many of the genes mentioned here have additional functions in vegetative growth and in spikelet development, but pleiotropy may obscure their role in any single process, hence the focus on a narrow slice of time. Specifically, I do not address the vegetative–reproductive transition and control of flowering time, which are thoroughly discussed elsewhere (e.g. Distelfeld et al., 2009; Lee and An, 2015; Doust et al., 2017; Woods et al., 2019), nor do I review the fundamental controls of IM size, which are also covered in recent reviews (e.g. Bommert and Whipple, 2018). The extensive data on floret structure, form, and function (e.g. Schrager-Lavelle et al., 2017; Shen et al., 2021) are also not covered here. Some of these topics are included in the related review by Richardson and Hake (2022), which focuses particularly on the incomparable data available from maize, and also recent species-focused reviews on rice (Li et al., 2021a, 2021b, 2021c) and Triticeae (Gao et al., 2019; Gauley and Boden, 2019; Sakuma and Schnurbusch, 2020). Protein-coding genes discussed in this review are listed in Supplemental Table S1.

Grass inflorescences are branched structures with branches producing spikelets

The grass family (Poaceae or Gramineae, both correct names) includes about 700 genera and 12,000 species (Kellogg, 2015; Soreng et al., 2017). The family is divided into 12 subfamilies, 3 of which (Anomochlooideae, Pharoideae, and Puelioideae) are successive sisters to the remainder of the family and together include only a handful of species (GPWG II, 2012; Saarela et al., 2018; Figure 1). The other nine subfamilies fall into two large clades that are strongly supported by phylogenetic data and are named by the




Figure 1 Phylogeny of the grass family summarized from GPWG II (2012); Soreng et al. (2017), and Saarela et al. (2018). Within the grasses, terminal taxa are subfamilies; representative crops are given where applicable.

acronym for the included subfamilies (Kellogg, 2015; Soreng et al., 2017). The BOP clade (with ~40% of the species in the family) includes Bambusoideae, Oryzoideae, and Pooideae, while the PACMAD clade (with ~60% of the species) includes Panicoideae, Aristidoideae, Chloridoideae, Micrairoideae, Arundinoideae, and Danthonioideae.

Grass inflorescences are distinctive. In all but the four species of Anomochlooideae (Judziewicz and Soderstrom, 1989), the flowers (florets) are borne in units known as spikelets (GPWG, 2001; Figure 2). Each spikelet consists of sterile bracts (glumes, generally two) and a spikelet axis bearing one or more florets. The number of florets per spikelet is generally fixed within a species or clade or varies within a narrow range. The flowers themselves are zygomorphic, with a large subtending bract (the lemma), in the axil of which is a conventional, if highly reduced, flower with an outer perianth (the palea), inner perianth (lodicules), stamens, and a gynoecium with a plumose stigma and single ovule. Florets are thus determinate structures and their formation marks an end to any further branching processes.

Grass inflorescences are often described as spikes, racemes, or panicles, borrowing terminology from dicots and non-grass monocots. However, because the terminal units are spikelets, which are themselves inflorescences, rather than flowers as in dicots, the grass inflorescence is in fact a compound structure, an inflorescence of spikes, and is therefore technically a synflorescence (Weberling, 1989; Vegetti and Weberling, 1996). I will use the more common term inflorescence here even though it is somewhat inaccurate.

In this review, I treat the grass family as a single genetic system (Bennetzen and Freeling, 1993), with insights coming from cross-species comparisons as much as from detailed studies in a single species. Data come largely from rice (*Oryza sativa*, tribe Oryzeae, subfamily Oryzoideae) and maize (*Zea mays*, tribe Andropogoneae, subfamily Panicoideae), barley, and wheat (*Hordeum vulgare* and *Triticum aestivum*, respectively, tribe Triticeae, subfamily Pooideae; Figures 1–3; gene names in Supplemental




Figure 2 Images and diagrams of spikelets of wheat (*Triticum turgidum* cv. “Kronos”), rice (*O. sativa*), barley (*H. vulgare*, only central spikelet diagrammed), and maize (*Z. mays*, tassel spikelet). Paleas not visible in photos of wheat and barley. Shaded ovals (yellow) indicate floral organs as shown in inset, upper right. Glumes indicated by thickened arcs (green); lemmas and paleas, black arcs; suppressed meristems, x. Scale bars, 0.5 mm. Distance between structures in diagrams is exaggerated for clarity. Image of barley spikelet reproduced from Komatsuda et al. (2007); copyright 2007 National Academy of Sciences. Scale bar lacking in original. an, anther; gl, glume; lo, lodicle; pa, palea; st, stigma.

Table S1), although I will also mention *Brachypodium distachyon* (tribe Brachypodieae, subfamily Pooideae), and green millet (*Setaria viridis*, tribe Paniceae, subfamily Panicoideae). No universal system of gene nomenclature exists for the grasses, so for consistency gene and protein names are written in all capital letters, with the gene names italicized. I do not try to distinguish orthographically between dominant and recessive alleles.

Inflorescence architecture is the outcome of a repeating series of developmental decisions

The pattern of inflorescence architecture is governed by the relative timing of shifts from branch-producing to

spikelet-producing meristems, what Kyozyuka (2014) has described as a “meristem phase change.” In all grasses after the transition of the SAM to reproductive activity, an IM or BM has only three possible fates: (1) it can function as a branch-producing meristem, with new BMs arising on its flanks; (2) it can form a spikelet (SM), or (3) it can cease to function, with cell division and growth coming to a halt. If the BM produces higher order BMs, these in turn have the same set of developmental options. Production of glumes marks the transition from a BM to an SM, and further development is canalized to produce one or more florets.

The inflorescence architecture of grasses is well documented and shows that the number of iterations of these developmental decisions varies among species and genera.

However, for any given species, the number of BMs produced by an IM or other BMs is fixed within a narrow range as is the number of SMs.

Inflorescence development may be modeled as a series of on–off switches, with shifting patterns of identity (Kellogg, 2000). In contrast, Prusinkiewicz et al. (2007) and Harder and Prusinkiewicz (2013) describe a model more similar to a rheostat with continually varying amounts of what they term “vegetativeness.” The models are not mutually exclusive, in that the switch model can be viewed as a simple version of the rheostat model. However, the latter model has never been elaborated formally to accommodate the complexities of grass inflorescence architecture.

The branching pattern of grass inflorescences is established early in development when the IM is still enclosed by the sheathing leaf bases. Elongation of inflorescence internodes occurs later (Patil et al., 2019; McKim, 2020; E. A. Kellogg, personal observation). Most research has focused on branching patterns because they determine the number of florets and hence the number of grains, and this review reflects that same bias, with few comments on elongation. Nonetheless, inflorescence elongation patterns may have fitness consequences in both natural and agricultural settings, for example, by determining whether the florets must pollinate themselves or can shed pollen away from the plant, or determining the propensity of the inflorescence stalk to break or be attacked by herbivores.

Whether the IM ultimately produces a spikelet is independent of whether the primary BMs terminate in spikelets or simply cease growing. The IM ceases producing BMs and ends as an undifferentiated dome or axis in maize (*Z. mays*), rice (*O. sativa*), signal grass (*Brachiaria decumbens*), and finger millet (*Eleusine coracana*; Figure 3), whereas it ultimately becomes an SM in wheat (*T. aestivum*), ryegrass (*Lolium* spp.), sorghum (*Sorghum bicolor*), and oats (*Avena sativa*; Butzin, 1979; Moncur, 1981; Liu et al., 2007; Reinheimer and Vegetti, 2008; Reinheimer et al., 2009, 2013; Kellogg et al., 2013). While presence of a terminal spikelet is consistent within a species or genus, it is labile in evolutionary time (Reinheimer and Vegetti, 2008; Reinheimer et al., 2013; Kellogg, 2015), suggesting it can be altered easily by natural selection but is generally not a plastic response to the environment.

Placement of bracts is governed by auxin and specifies branching patterns

The role of auxin

Branches in all plants form from meristems in the axils of leaves or bracts, which in turn are controlled by local auxin maxima leading to lateral organ production (Reinhardt et al., 2003; Smith et al., 2006). In rice and maize, auxin accumulates in all BMs of the inflorescence, as well as in the primordia of glumes and other floral organs (Yang et al., 2017). Auxin biosynthesis, transport, and signal transduction have been recently reviewed by Matthes et al. (2019), who provide detailed information on the molecular evolution and




Figure 3 Inflorescence diagrams of rice (*O. sativa*), barley (*H. vulgare*), wheat (*T. aestivum*), and maize (*Z. mays*). Spikelets indicated by ovals (green); suppressed bracts, arcs below the ovals (orange). IMs lacking a terminal flower indicated by asterisks (magenta).

expression of auxin-related genes in rice and maize and compare auxin biology in grasses with that in Arabidopsis. Bract and axillary branch formation require auxin biosynthesis, as shown by disruption of the auxin biosynthetic genes in maize *VANISHING TASSEL2* and *SPARSE INFLORESCENCE1* and their rice orthologs *OsTAR2/OSFISHBONE* and *OsYUC1* (Gallavotti et al., 2008; Phillips et al., 2011; Yoshikawa et al., 2014; Matthes et al., 2019). All are expressed in axillary meristems of the inflorescence.

Auxin flux through the epidermis converges on small regions of the IM, a process regulated by the auxin influx carrier *ZmAUX1* (*SvAUX1* in *S. viridis* (Zhu et al., 2021a, 2021b) and the auxin efflux carrier *SISTER OF PINFORMED1* (*SoPIN1/ZmPIN1D*; O'Connor et al., 2014; Matthes et al., 2019; Figure 4). Loss-of-function mutations in *SvAUX1/SPARSE PANICLE1* reduce all orders of branching in the inflorescence, whether primary, secondary, tertiary, or higher, although the effect in maize is less striking than in *S. viridis* (Huang et al., 2017; Zhu et al., 2021a, 2021b). Vein patterning is controlled by *PIN-FORMED1a* (*PIN1a*) and *PIN1b*, which move auxin away from the local maxima and establish the position of vascular tissue (O'Connor et al., 2014). *SoPIN1/ZmPIN1D* is shared by all angiosperms except Brassicaceae, where the gene appears to have been lost




Figure 4 BA1/LAX1 (boundary domain), RA2 (axillary meristem), and TSH4 (suppressed bract) regulatory networks. Subnetworks in boxes are widely conserved among the grasses. ^aOrtholog in rice not involved in regulating LAX1/LAX2. ^bNo data on orthologs in other grasses. ^cRA2/HvRA2 are expressed adjacent to the meristem but not overlapping with BA1 + BA2. ^dGene absent in rice and barley genomes. Dashed lines indicate regulatory connection inferred from gene expression and mutant phenotypes, following Yao et al. (2019). For genes with different names in different species, the first name is the maize gene name (black), second is rice (magenta), and third is barley (cyan).

(O'Connor et al., 2014; Matthes et al., 2019). In rice, mutations in *OsPIN1a* and *OsPIN1b* primarily affect the root system, although *PIN1a* mutants and the double mutant *PIN1a PIN1b* exhibit increased branch angle in the inflorescence (Li et al., 2019a, 2019b). The single mutants *PIN1c* and *PIN1d* had no effect on the inflorescence, but the double *PIN1c PIN1d* mutant abolished all inflorescence branching (Li et al., 2019a, 2019b). The wheat TaPIN1 proteins also affect spikelet number and grain number per inflorescence (Yao et al., 2021).

ZmPIN1a is phosphorylated by BARREN INFLORESCENCE2 (BIF2), a homolog of Arabidopsis PINOID (McSteen et al., 2007; Skirpan et al., 2008). Without BIF2, branches do not initiate. In rice, mutations of OsPID had no effect on inflorescence branching although they affected floral organ development (Xu et al., 2019).

The auxin/indole-acetic acid (Aux/IAA) proteins form a large complex family of proteins in the grasses (as in other plants), but appropriate genetic studies are largely lacking. Few AUX/IAA mutants are known to have clear effects on inflorescence branching (Matthes et al., 2019). Two such genes in maize, *BIF1/ZmIAA27* and *BIF4/ZmIAA20*, are expressed in the IM and in the central zone of axillary

meristems (Galli et al., 2015). BIF1 and BIF4 interact with maize activating auxin response factors (ARFs) and appear to regulate the placement of axillary meristems and to repress their formation elsewhere.

Phyllotaxis, a read-out of hormonal signals

In most grasses and their outgroups, the IM produces bracts and their axillary primary branches in a spiral, a shift from the distichous phyllotaxis of the vegetative shoot (Kellogg et al., 2013; Bartlett and Thompson, 2014). Although spiral inflorescence phyllotaxis is apparently ancestral in the grasses, all members of Pooideae except the early diverging genus *Brachyelytrum* (Kellogg et al., 2013), some Danthonioideae and many Andropogoneae (Panicoideae) maintain two-ranked bracts and branches in the inflorescence. Also in Pooideae, some inflorescences are two-ranked but with a divergence angle $< 180^\circ$, a pattern that is phenotypically similar to rice inflorescences with mutations in *ABERRANT PANICLE ORGANIZATION1* (*AP01*; Ikeda et al., 2005, 2007). In such inflorescences, the placement of the bracts does not follow the placement of the leaves (Kellogg et al., 2013). Truly distichous inflorescences appear in the

crown Pooideae (tribes Brachypodieae, Triticeae, Bromeae, and Poeae).

Inflorescence bracts are suppressed; axillary meristem growth is continuous

Bract suppression

In nearly all grasses, inflorescence bracts are suppressed and are apparent only in early development as narrow ridges (see e.g. Moncur, 1981; Kellogg et al., 2013), a pattern distinct from most other angiosperms and grass outgroups in which inflorescence bracts expand. Conversely, axillary branches in grass inflorescences grow immediately upon initiation (Kyoizuka 2014; Li et al., 2019a, 2019b). Inflorescence bract suppression contrasts with that of vegetative growth, in which the leaves expand but the axillary meristems generally fail to grow out immediately (Kyoizuka 2014). However, bracts do expand in some parts of the inflorescence of Bambusoideae, many Andropogoneae, and possibly in *Anomochloa*, although the morphology of the latter is complex and hard to interpret (Judziewicz and Soderstrom, 1989; Judziewicz et al., 1999; Sajo et al., 2012).

The proteins TASSELSEATH1 (TSH1) and TSH4 repress inflorescence bracts; when both are mutated, bracts expand and axillary meristems are reduced or entirely absent (Whipple et al., 2010; Xiao et al., 2021). TSH1 is a GATA-domain zinc finger transcription factor orthologous to NECKLEAF1 (NL1) in rice and THIRD OUTER GLUME (TRD) of barley (Houston et al., 2012), which have similar mutant phenotypes and together are the founding members of the NL1/TSH1/TRD (NTT) family of proteins. TSH4 is a transcription factor in the SQUAMOSA-PROMOTER BINDING (SBP)-LIKE (SPL) family; it is orthologous to ZmSBP23 and the two maize proteins are co-orthologous to OsSP17 (Wei et al., 2018).

Both *TSH4* and *TSH1* are expressed in the cryptic bract of inflorescence branches and act synergistically, with *TSH4* upstream of *TSH1* (Xiao et al., 2021; Figure 4). Together they regulate the expression of *LIGULELESS2* (*LG2*), encoding a basic leucine zipper protein that regulates outgrowth of long branches among many other phenotypes (Walsh and Freeling, 2002). *TSH1* is co-expressed with *ZmYABBY15*, a gene expected to be expressed in all leaf-like organs, confirming that the minute structure is indeed a bract. Based on gene expression and mutant phenotypes, Xiao et al. (2021) suggest that the maize SPL proteins UNBRANCHED2 (*UB2*) and *UB3* may also help regulate *TSH1* expression. *UB2* and *UB3* are co-orthologous to OsSPL14 (Chuck et al., 2014) and may control allocation of cells to lateral organs in general.

TSH1 and the NTT proteins NTT-like1 and NTT-like2 all share a HAN domain with the dicot proteins related to HANABA TARANU (*HAN*; Whipple et al., 2010), although the NTT proteins are all more closely related to each other than any of them is to the dicot proteins. Despite the sequence similarities among grass (NTT) and dicot *HAN* proteins, *HAN* does not regulate bract development, even

though bracts are suppressed in Brassicaceae. Thus, bract suppression in the grasses must proceed by a mechanism quite different from that of *Arabidopsis* (Whipple et al., 2010). NTT-like genes are expressed in boundary regions in outgroups of the grasses, but early in grass evolution *TSH1* acquired SBP binding sites in its promoter, bringing it under the control of TSH4 and causing bract expression (Xiao et al., 2021). In this position, TSH1 retains its presumed ancestral developmental role of suppressing cell division and growth, as well as a possible role in signaling.

Axillary meristem growth: the BA1/LAX1 network

Signaling centers adjacent to axillary meristems appear to specify which cells develop as part of the bract, which become part of the meristem, and which cease dividing entirely and form a boundary. Regulatory proteins in these centers likely also activate mobile factors that then move into the meristem (Whipple, 2017). Such regulatory proteins include a conserved basic helix–loop–helix transcription factor, LAX PANICLE1 (*LAX1* in rice, orthologous to BARREN STALK1, *BA1*, in maize) that is required for axillary branch formation (Komatsu et al., 2003a, 2003b; Gallavotti et al., 2004; Figure 4). In all grasses that have been investigated, the underlying genes are expressed in the boundary between the axillary meristem and the axis that bears it (Komatsu et al., 2003a, 2003b; Gallavotti et al., 2004; Oikawa and Kyoizuka, 2009; Woods et al., 2011). *BA1/LAX1* mutations do not affect bract formation so are specific to the region just adaxial to the meristem. The auxin transport protein BIF2 (*OsBIF2* in rice) physically interacts with and phosphorylates *BA1* (Skirpan et al., 2008).

BA1/LAX1 interacts directly with *BA2/LAX2*, a nuclear-localized protein that is conserved in grasses (Figure 4). The expression domains and mutant phenotypes of *BA1/LAX1* and *BA2/LAX2* overlap, further supporting their involvement in the same pathway (Tabuchi et al., 2011; Yao et al., 2019). *BA1/LAX1* is expressed in *BA2/LAX2* mutants and vice versa, indicating that the proteins do not regulate each other's transcription (Tabuchi et al., 2011; Yao et al., 2019).

Other regulators of *BA1/LAX1* have been identified in maize and rice but have been investigated in only one of the two species. For example, MONOCULM1 (*MOC1*) in rice, a transcription factor with a GRAS domain similar to that of LATERAL SUPPRESSOR in *Arabidopsis*, positively regulates *LAX1* (Li et al., 2003). *MOC1* mutants have fewer branches in the inflorescence (Li et al., 2003), and the *LAX2 MOC1* double mutant lacks branches entirely (Tabuchi et al., 2011). Mutants in *MOC1* orthologs have not been described in other grasses.

Likewise, BARREN STALK FASTIGIATE1 (*BAF1*) in maize, an AT-hook domain DNA binding protein with a plant-specific Plant and Prokaryote Conserved domain, positively regulates *BA1* although the two do not interact directly (Gallavotti et al., 2011). In *BAF1* mutants, axillary meristems are fused to the axis, but bracts are unaffected (Gallavotti et al., 2011). *BAF1* may also regulate axillary meristems directly, bypassing *BA1* (Matthes et al., 2019). The *BAF1*

ortholog in rice affects floral development and is known as *PALEALESS1* or *DEPRESSED PALEA1* (*DPI1*; Jin et al., 2011). However, a branching phenotype is not reported in rice, nor is a floral phenotype in maize.

Expression of *RAMOSA2* (*RA2*, orthologous to barley *HvRA2* (*VRS4*) and rice *OsRA2*) marks the position of primary and secondary BMs in maize, sorghum, rice, and barley (Bortiri et al., 2006; Figure 4). *RA2* is a Lateral Organ Boundaries (LOBs)-domain transcription factor that is conserved in grasses (Bortiri et al., 2006; Koppolu et al., 2013), and has a grass-specific sequence upstream of the LOB domain (Koppolu et al., 2013). In normal development, the IM in both maize and barley produces short lateral branches each of which produces only two (maize) or three (barley) spikelets (Figure 3); in the maize and barley literature, the meristems producing these short branches are known as a spikelet pair meristem and a triple mound, respectively. Mutations in *RA2* and *HvRA2* permit the short branches to continue growth, leading to a branch with unpaired spikelets in maize (Bortiri et al., 2006), and a branch-like central spikelet and fertile lateral spikelets in barley (Koppolu et al., 2013). This continued growth reflects a delay in terminal spikelet formation, also described as loss of determinacy. In contrast, downregulation of *OsRA2* did not affect branching but pedicel length increased, indicating that the normal function of the protein in rice is to prevent growth of specific tissues, but possibly not inflorescence branches (Lu et al., 2017). Branch length was not affected, although overexpression of *OsRA2* reduced the number of secondary branches.

Opposing regulatory gradients of *miR156*-SPL control branching

Some developmental decisions can be described as transitions and gradients, with the gradients often running in opposition to each other. microRNAs and their targets have become well known for setting up such opposing gradients. For example, the microRNA *miR156* is upregulated by SPL proteins; it then cleaves the corresponding SPL transcript in a negative feedback loop, a process initially elucidated in vegetative to reproductive phase change in Arabidopsis and maize (Chuck et al., 2007a, 2007b; Poethig, 2009; Wu et al., 2009). In Arabidopsis, as *miR156* expression decreases, *SPL3* expression goes up, increasing expression of *LFY*, *AP1*, and *FUL* (Yamaguchi et al., 2009).

In grasses, several SPL proteins and their regulatory microRNAs control the transition from branching to spikelet production. The rice genome includes 19 *OsSPL* loci, 11 of which could be targets of *miR156* based on sequence comparisons (Xie et al., 2006; Yang et al., 2008); comparable numbers in maize are 30 and 18, respectively. Many rice SPL loci were discovered initially as quantitative trait loci in studies aiming to improve grain number; because of that history, many have been named more than once in the literature. A full list of alternative gene names is in Supplemental Table S1. Among the loci with *miR156* binding sequences are

OsSPL6 (*ZmSBP6*, 17), *OsSPL8* (*ZmLG1*), *OsSPL13* (*ZmSBP13*, 29), and *OsSPL16*, *OsSPL18* (the latter two co-orthologous to *TEOSINTE GLUME ARCHITECTURE1* (*TGA1*), *NEIGHBOR OF TGA1* (*NOT1*), and *ZmSBP5*; Wei et al., 2018). Other highly expressed *OsSPL* loci include *OsSPL7*, *OsSPL14*, and *OsSPL17* (Wang et al., 2015).

OsSPL14 (orthologous to maize *UB2* and *UB3*) has received particular attention (Jiao et al., 2010; Miura et al., 2010). Increased transcription of *OsSPL14* leads to more primary inflorescence branches (Huang et al., 2016), and heterozygotes were strongly over-dominant for yield (reflecting higher ultimate numbers of spikelets; Figure 5A). Mutation of the *miR156* binding site also increased *OsSPL14* expression and yield (Jiao et al., 2010). Overexpression of *OsSPL14* or inhibition of *miR156* both led to early transition from BMs to SMs. Consistent with this interpretation, expression of *FRIZZY PANICLE1* (*FZP1*; a spikelet marker, see below) was higher and *FZP* was expressed in meristems that might otherwise have produced branches (Wang et al., 2015). Mutant phenotypes of *OsSPL14* and *OsSPL17* are similar, with double mutants (RNAi) showing enhanced effects.




Figure 5 A, *miR156*-SPL-*miR172*-AP2-like regulatory networks. B, Developmental window showing the transition from axillary BMs to spikelet formation, with major genes marking each stage and opposing gradients of microRNAs. Species- and clade-specific inflorescence morphology is influenced by the developmental timing of the transition; shorter time causes a faster transition to glume production which in turn leads to fewer branches and vice versa. In rice, transition time appears to vary continuously across the inflorescence; in maize, transition time is bimodal (long and short, but nothing in between); in barley, transition time is unimodal, only short. Portions of the figure are redrawn from Wang et al., (2015, Supplemental Figure S16).

The complex of APO1, an F-box protein and APO2 (RFL), a homolog of Arabidopsis LEAFY, also delays the transition from BMs to producing glumes (Ikeda et al., 2005; Ikeda-Kawakatsu et al., 2009, 2012). When the LFY homologs ZFL1 and ZFL2 are mutated in maize, the transition to normal tassel branches is also delayed and axillary meristems develop in husk leaves of the ear (Bomblies et al., 2003).

The direct interaction of APO1 and APO2 appears to be conserved in plants, having been demonstrated in Arabidopsis (orthologs UNUSUAL FLORAL ORGANS and LFY, respectively), rice, and barley (Chae et al., 2008; Kyoizuka, 2014; Selva et al., 2021). However, the two proteins delay the transition to spikelet formation in rice and maize in whereas they promote flower formation in Arabidopsis (Kyoizuka, 2014). Their role in barley is still different, in that disruption of HvLFY does not affect inflorescence architecture nor does it affect expression of *APETALA1/FRUITFULL* (*FUL*)-like (*FUL*-like) genes, as might be expected if it affected spikelet formation (Selva et al., 2021). WFL (ortholog of LFY in wheat) is expressed in the bracts (lower ridge) below the spikelets, but not in the spikelets themselves (Shitsukawa et al., 2006); its mutant phenotype is unknown.

Control of secondary and higher order branches is species- and position specific

Both the number and morphology of secondary and higher order branches vary among cereal grasses (Figure 3), among genera within a tribe (e.g. Cynodonteae; Pilatti et al., 2019), and among species within a genus (e.g. *Setaria*; Doust and Kellogg, 2002). The range of natural variation suggests that genetic control of secondary branches may be partially independent of the control of primary branches, a hypothesis supported by genetic data.

Secondary branching in rice

Primary and secondary branches are controlled independently in rice, even though all branches are morphologically similar (Harrop et al., 2019; Bai et al., 2021). For example, double mutants of *LAX1LAX2* (described above), have no visible defect in primary branches but lack secondary branches and spikelets altogether (Tabuchi et al., 2011). Allelic variation has been explored extensively in *FZP1* (see also below), an AP2/EREBP transcription factor (Bai et al., 2017; Fujishiro et al., 2018; Huang et al., 2018; Wang et al., 2020), where mutations in the promoter affect the binding of transcription factors and thereby expression levels. An allele of *FZP1* originally known as *CONTROL OF SECONDARY BRANCH1* acts particularly on secondary branches (Huang et al., 2018). A small deletion in the promoter of *FZP1* reduces binding of the ARF OsARF6 and reduces *FZP* expression, leading to increased cell division and more secondary branches, but no changes in primaries.

In addition, *FZP1* interacts with and is degraded by NARROW LEAF 1 (*NAL1*), a serine/cysteine protease. Downregulation of *FZP1* along with upregulation of *NAL1* improved yield in rice (Huang et al., 2018), again by

increasing secondary branches. *NAL1* is expressed throughout the plant, particularly in vascular tissues (Qi et al., 2008), whereas *FZP1* is expressed only in the inflorescence.

Knockout of *OsSPL18* significantly reduces the number of secondary branches, and *OsSPL18* is itself cleaved by *OsmiR156k* (Yuan et al., 2019). *OsSPL18* binds to the promoter of *DENSE and ERECT PANICLE1* (a G-protein γ subunit; Xing and Zhang, 2010; Liu et al., 2021) and activates it (Yuan et al., 2019), thereby increasing cell numbers. Mutations in *OsSPL9*, the gene underlying the mutant *LESS GRAIN NUMBERS5*, exhibited less than half the number of secondary branches as wild-type *indica* lines, although primary branch number was unaffected (Hu et al., 2021).

Overexpression of *RICE CENTRORADIALIS1* and 2 (*RCN1*, *RCN2*), homologs of *CENTRORADIALIS/TERMINAL FLOWER1*, led to increased panicle branching in rice (Nakagawa et al., 2002; Wang et al., 2015), indicating that the primary function of the RCNs is to reduce branching, perhaps by accelerating the transition to spikelet formation. Overexpression of *RCN* rescued the effects of *OsSPL14* and *OsSPL17* RNAi lines on secondary branches but did not affect primary branches (Wang et al., 2015).

OsMADS34/PANICLE PHYTOMER2 (*PAP2*) in rice also controls the relative numbers of primary and secondary branches (Gao et al., 2010; Kobayashi et al., 2010) with the normal function to reduce numbers of primary branches. The effect on secondary branching is unclear, with some mutations leading to more secondary branches and hence spikelets (Kobayashi et al., 2010), while others reported mutations lead to fewer (Gao et al., 2010).

The regulatory networks controlling secondary branch formation in rice may be relevant in other species with open branching inflorescences such as the closely related genus *Zizania* (North American wild rice) or the distantly related genera *Panicum* (switchgrass) or *Megathyrsus* (guinea grass). However, other species have distinct architecture and are hard to compare to rice (Figure 3).

Secondary branching in maize: spikelet pairs

Maize produces long and short inflorescence branches. In the tassel, the first-formed primary branches are long, whereas later ones are short, producing exactly two spikelets (spikelet pairs; Figure 3). In these, one spikelet is lateral (i.e. a secondary branch) and the other is terminal. Primary branches in the ear are also short (spikelet pairs) as are secondary branches in the tassel.

The maize branch regulator *RA2* is genetically upstream of *RA1*, which has a similar mutant phenotype in which spikelet pairs are converted to longer branches, often with single spikelets (Vollbrecht et al., 2005; Bortiri et al., 2006; Figure 4). *RA1* is a C2H2 zinc-finger transcription factor containing two Ethylene-responsive element binding factor-associated Amphiphilic Repression (EAR) domains (Vollbrecht et al., 2005). *RA1* interacts directly with *RAMOSA ENHANCER LOCUS1* (*REL1*), orthologous to *ABERRANT SPIKELET AND PANICLE1* (*ASP1*) in rice (Gallavotti et al., 2010; Tanaka et al., 2013). *REL1/ASP1* is

a protein with an AT-hook domain similar to TOPLESS in *Arabidopsis* and is thought to be a transcriptional co-repressor. However, because rice lacks a RA1 locus (Vollbrecht et al., 2005), the interactors of ASP1 are unclear.

RA3, a trehalose-6-phosphate phosphatase (TPP) in maize, is genetically independent of RA2 although it also affects the short branch/spikelet pair meristem (Satoh-Nagasawa et al., 2006; Figure 4). TPP may link sugar metabolism to signaling, since other TPPs repress *SUCROSE-NON-FERMENTING1-RELATED KINASE1* and also *miR156* which in turn negatively regulates SPL proteins (Eveland and Jackson, 2012; Tsai and Gazzarrini, 2014). However, RA3 co-localizes with RNA POLYMERASE II in nuclear speckles (Demesa-Arevalo et al., 2021), and an RA3 construct lacking phosphatase activity will still complement the *ra3* mutant (Claeys et al., 2019). Together these observations suggest a transcriptional regulatory role for RA3 separate from its role as an enzyme.

Secondary branching in barley: triplets of spikelets

RA1 and RA3 are absent from genomes in the BOP clade, so rice, barley, wheat, and other related species must have distinct pathways regulating higher order branching (Vollbrecht et al., 2005; Doust, 2007; Kellogg, 2007). In barley, the primary branches are short and terminate in a spikelet, but before terminating they produce exactly two lateral (secondary branch) spikelets. While this complex of three spikelets is formed from a triple meristem, in barley relatives such as *Elymus* and *Leymus* the number of secondary spikelets varies from one to three depending on the species (POWO, 2021).

The controls of secondary (lateral) branching in barley reflect a complex network involving branching, glume formation, and floral organ development (Gauley and Boden, 2019). Branching itself (i.e. formation of the laterals) is governed by HvRA2 but proteins regulated by HvRA2 differ from those of maize RA2, which is unsurprising given the lack of RA1 and RA3 in barley (Figure 4). As in maize, a TPP protein is genetically downstream of HvRA2, but the barley TPP protein is HvSRA, which is not orthologous to RA3 but rather belongs to the SISTER OF RA3 (SRA) clade of TPPs that is conserved in grasses. SRA is apparently not involved in inflorescence development in maize (Satoh-Nagasawa et al., 2006). HvRA2 upregulates *VRS1/HOX1*, a homeodomain leucine zipper transcription factor that is the result of a gene duplication specific to Triticeae (Komatsuda et al., 2007; Koppolu et al., 2013; Sakuma et al., 2019). The HOX1/HOX2 clade is in turn sister to orthologs of GRASSY TILLERS1 in maize (Whipple et al., 2011). Recent work on barley MADS-box transcription factors also suggests they regulate inflorescence branching, possibly in response to temperature, in addition to their expected function in floral organ identity (Kuijter et al., 2021; Li et al., 2021a, 2021b, 2021c). Thus, despite the conserved LOB-domain transcription factors (RA2 and HvRA2) and the involvement of a TPP protein, the controls of spikelet pairs and lateral branches in Triticeae differ from those in other grasses.

Spikelet bracts (glumes) expand, axillary bud growth is suppressed

Glume production is common but not a necessary marker of a transition to floret production

The shift from a BM with suppressed bracts and active axillary meristems to a spikelet-producing meristem is clear in many grasses, with the SM producing exactly two macroscopic bracts (glumes) with suppressed axillary meristems, followed by one or more large bracts (lemmas) subtending FMs. However, that transition may be protracted, with some species producing more than two glume-like structures. For example, the glumes in rice are tiny and known as rudimentary glumes (Figure 2). Distal to the glumes are two structures in the position of florets that also fail to produce axillary FMs. Although many lines of evidence support the inference that these are sterile lemmas, they are expanded bracts without an axillary meristem so share some characteristics with glumes. Mutations in *G1/LONG SLENDER LEMMA1* (Yoshida et al., 2009; Yang et al., 2020) shift the size and cellular morphology of the sterile lemmas to look more like true lemmas, whereas mutations in other genes lead to stronger similarity between the sterile lemmas and rudimentary (true) glumes (summarized by Ren et al., 2018; Xu et al., 2020). However, none of the mutants leads to production of an axillary FM, as would be expected if the sterile lemmas were fully converted to true lemmas. Other species (e.g. *Chasmanthium*) have multiple sterile lemmas, whereas many Bambusoideae bear pseudo-spikelets, which are subtended by glume-like structures with axillary meristems that themselves produce spikelets, somewhat reminiscent of FZP mutants (see below).

Conversely, the absence of glumes is well documented in some species of grasses, including many members of Oryzae (some of which also lack sterile lemmas), as well as *Nardus* and *Lygeum* in the tribe Nardeae (subfamily Pooideae), tribe Orcuttieae (Chloridoideae), *Pirenia* (Bambusoideae), and others (Kellogg, 2015).

The relative timing of glume production determines the overall architecture of the inflorescence and thereby the potential for seed production. In all grasses studied, meristems in the axils of glumes are suppressed by BRANCHED SILKLESS1 (BD1; maize)/FZP1; rice), orthologs of which have been characterized in *B. distachyon*, barley, and wheat (Chuck et al., 2002; Komatsu et al., 2003; Zhu et al., 2003; Derbyshire and Byrne, 2013; Dobrovolskaya et al., 2015; Poursarebani et al., 2015). Meristems form in the axils of glumes in BD1/FZP1 mutants, and these axillary meristems each produce glumes with axillary meristems. FZP is thus central to the transition from BM with suppressed bracts to SM with suppressed axillary meristems. The broad phylogenetic distribution of these systems indicates that the function of FZP1 orthologs is likely conserved among all spikelet-bearing grasses.

BD1/FZP1 is a transcription factor with a single AP2 domain, and part of the AP2/ERF clade similar to PUCHI in *Arabidopsis* (Chandler, 2018), which also specifies axillary meristems in the inflorescence (Karim et al., 2009). Unlike

BD1/FZP1, PUCHI appears to promote axillary meristem (flower) growth rather than suppress it. The sequence of the BD1/FZP AP2 domain is conserved across grasses. Two FZP1 homologs have been identified in the genome of *Pharus latifolius*, a member of the subfamily Pharoideae that is sister to all other spikelet-bearing grasses (Ma et al., 2021; Figure 1), although only one gene is expressed in young inflorescences.

Transition from producing suppressed bracts to producing glumes controls numbers of branches, particularly secondaries

Within the inflorescence and along each branch, gradients of gene regulation, particularly via *miR172* and *euAP2*-like genes, control the transition from BM production (suppressed bracts) to SM production (expanded bracts; Wang et al., 2015; Figure 5B). The relative speed of this transition controls the architecture of the inflorescence (Kyojuka, 2014).

euAP2-like genes have two AP2 domains (Kim et al., 2006), rather than the single domain found in FZP. In addition, nearly all genes in this group also have *miR172* binding sites (Seetharam et al., 2021) and *miR172* expression opposes expression of AP2-like genes (Figure 5A). The AP2-like-*miR172* interaction has been investigated for its role in specifying the number of florets per spikelet (e.g. Chuck et al., 2007, 2007b, 2008; Zhu et al., 2009), but another important role of AP2-like-*miR172* is to delay production of glumes, thereby prolonging branching.

In single and double mutants of the rice AP2-like genes *INDETERMINATE SPIKELET1* (*OsIDS1*) and *SUPERNUMERARY BRACT* (*SNB*), the IM and BMs were converted precociously to spikelets, leading to fewer branches (both primary and secondary), with the number varying in a dose-dependent manner (Lee and An, 2012). Mutant spikelets had extra rudimentary glumes, indicating that the meristem had made a transition from producing suppressed bracts (as in a branch) to producing glumes (as in a spikelet), but had failed in the subsequent transition to FM production (Lee and An, 2012). Consistent with this interpretation, FZP expression appeared earlier in BMs of the mutants than in wild-type. Overexpression of *miR172* in rice produced a phenotype similar to that of the *OsIDS1* *SNB* double mutant. Mutations in the orthologous genes in maize (*IDS1* and *SISTER OF IDS* [*SIDS*]) showed similar phenotypes, with fewer branches and extra glumes (Chuck et al., 2008).

In wheat, the *IDS* ortholog is the domestication gene *Q* (Seetharam et al., 2021), which is also regulated by *miR172* (Debernardi et al., 2017). The transition to forming glumes is particularly obvious in wheat because the glumes have prominent keels, shorter awns, and more sclerenchyma than lemmas. Reduction of *miR172* led to higher levels of *Q* (AP2-5) and greater similarity between glumes and lemmas. Conversely high levels of *miR172* and loss-of-function AP2-5 led to sterile lemmas. In the lowermost spikelets, the transition between glumes and lemmas appeared particularly malleable, such that more *miR172* and less AP2-5 could lead to glume-like organs in the position of lemmas (i.e. sterile lemmas).

Some SPL proteins, such as *OsSPL7* and *OsSPL14*, directly regulate *miR172* in rice and accelerate the transition to producing glumes (Wang et al., 2015; Figure 5A). The mutant phenotype caused by overexpression of either SPL locus was returned to normal by knockdown of *miR172*. Overexpression of *RCN1* and *RCN2* also led to increased panicle branching in rice by delaying the transition to SMs (Nakagawa et al., 2002; Wang et al., 2015).

TAWAWA1 is an ALOG protein that controls the timing of IM degeneration in rice and also the transition from BM to spikelet formation (Yoshida et al., 2013). Kyojuka (2014) has proposed that TAW1 is central to meristem maintenance in the IM and BMs, with lower levels leading to early IM abortion and accelerated transition from BMs to spikelet formation. TAW1 regulates *SHORT VEGETATIVE PHASE* (*SVP*) genes, which encode MADS-box transcription factors (Arora et al., 2007; Lee et al., 2007).

SM identity reconsidered

The existence of axillary signaling centers and gradients of developmental signals suggests that SM identity may be achieved by the confluence of several gene expression patterns that, when overlapping, produce the stereotypical grass spikelet. However, such patterns could also activate SM identity genes that are both necessary and sufficient to specify a structure as a spikelet. SEP-like and FUL-like MADS-box genes are good candidates for SM identity controls (Bommert and Whipple, 2018) as are the SPL proteins TGA1 and NOT1 (Preston et al., 2012).

MIKC-type MADS-box genes are well known as homeotic selector genes and some aspects of their function, particularly B-class (generally inner perianth and stamen expression patterns) and C-class (generally stamen and carpel expression) are conserved between dicots and grasses (Bommert et al., 2005). In contrast, the A-class function, originally thought to specify sepal identity and attributed to AP1, has been elusive (Litt and Irish, 2003; Litt, 2007). Grasses lack an ortholog of the dicot AP1 and instead have three loci that are more closely related to FUL in dicots (Preston and Kellogg, 2006, 2007). The three proteins, VERNALIZATION1 (*VRN1*; unrelated to the Arabidopsis protein of the same name), FUL2, and FUL3, affect plant height and flowering time in wheat, rice, Brachypodium, and Setaria (Yan et al., 2003; Kobayashi et al., 2012; Ream et al., 2014; Li et al., 2016; Woods et al., 2016; Li et al., 2019a, 2019b; Yang et al., 2021).

VRN1 and *FUL2* are expressed throughout the spikelet (glumes plus florets) in *Lolium* (ryegrass), *Triticum*, *Hordeum*, *Avena*, and *Setaria* (Gocal et al., 2001; Preston and Kellogg, 2008; Preston et al., 2009; Alonso-Peral et al., 2011; Yang et al., 2021), as well as being expressed in the IM and BM. Knockout of *VRN1FUL2* or *VRN1FUL2FUL3* in both genomes of tetraploid wheat specifically affected SM identity, consistent with their expression patterns (Li et al., 2019a, 2019b). In the mutants, the lower ridge expanded to form a leaf and the spikelet (the sole product of a primary branch) was replaced by a leafy tiller-like structure. Thus the ability of

the IM to position and form the subtending bract is not compromised but spikelet identity is disrupted. Having either VRN1 or FUL2 is enough to make a terminal spikelet and repress the wheat homologs of RCN; FUL3 controls timing of (accelerates) terminal flower production (Li et al., 2019a, 2019b). The balance between the FUL-like proteins and SVP proteins determines whether spikelets form normally in wheat, or whether they develop into tiller-like branches (Li et al., 2021a, 2021b, 2021c).

The rice proteins OsMADS5 and OsMADS34 (=PANICLE PHYTOMER2) add another layer of regulation. Like wheat VRN1 FUL2 FUL3 null triple mutants, the rice OsMADS14 OsMADS15 OsMADS18 PAP2 quadruple mutant replaces primary branches with vegetative tiller-like structures (Kobayashi et al., 2012; Li et al., 2019a, 2019b). In rice, OsMADS5 and OsMADS34 directly regulate RCN4 and accelerate the transition to spikelet production. Double mutants of OsMADS5 OsMADS34 or OsMADS34 RCN4 produce more branches, including secondary, tertiary, and even quaternary branches (Zhu et al., 2021a, 2021b). OsMADS34 promoters also contain SPL binding motifs, and OsMADS34 is directly regulated by OsSPL14 (Wang et al., 2015).

Mutations in OsMADS34 have no effect on rudimentary glumes, although the gene is expressed there; sterile lemmas in the mutants are morphologically similar to true lemmas but still do not produce axillary FMs (Gao et al., 2010). LACKING RUDIMENTARY GLUME 1 (LRG1) is also involved in glume and sterile lemma identity. In an unexpected example of regulatory convergence, LRG1 is a C2H2 transcription factor similar (although not orthologous) to RA1, with similar EAR repression domains and interactions with a TOPLESS-like protein (Xu et al., 2020). A full discussion of rice spikelet morphology is beyond the scope of this paper but will be interesting to pursue in the future.

Lee and An (2015) noted that expression of FUL-like MADS-box genes was unaffected in SNB OsIDS double mutants. One interpretation is that SNB and OsIDS are needed to establish the domain within which the FUL-like proteins can specify spikelet identity. Such an interpretation awaits additional data.

The SPL protein TGA1 acquired its current expression domain in the spikelet-bearing grasses (Preston et al., 2012) and is another candidate for conferring spikelet identity. In all grasses examined, it is expressed in the florets and both glumes. However, it is expressed only in the flower (not the floral bracts) of the grass outgroup *Joinvillea ascendens*. Thus, the grass expression pattern represents an expansion of floral control to encompass the bracts. Regulation of TGA expression has not been explored, although the *miR156* binding site is conserved among grasses and their outgroups (Preston et al., 2012).

Summary: conservation and diversity

The controls of inflorescence architecture are strikingly similar among many grasses (Figure 5B). Auxin transport and signaling use orthologous proteins retaining similar

biochemical functions, interactions, and developmental roles in most species. Likewise, conserved mechanisms specify the position and development of suppressed bracts via SBP proteins such as OsSPL18 and TSH1. BA1 and BA2 and their regulators and targets also appear conserved in positioning and delimiting axillary meristems. Spikelets are marked by formation of glumes; suppression of their axillary meristems is controlled by FZP/BD. Timing of transitions from IM to BM to SM is controlled by opposing gradients of *miR156-SPL-miR172-AP2*-like gene expression. This unifying mechanistic picture offers insights that may be applicable to less well-studied crops, as well as wild grasses.

Despite this broad similarity, many other mutant phenotypes have been observed only in a single species; it is unclear whether such gene functions are indeed phylogenetically restricted or if data on other species are simply lacking. For example, DP1 is the rice homolog of BAF1 and may have a different developmental role; however, the requisite data are not available. Allelic variation in FZP has been dissected carefully in studies in rice attempting to maximize yield, but no comparable data are available for BD1 in maize. Likewise, sets of genes control secondary branches (products of the primary BM) independent of primary branches (products of the IM) in rice, indicating that these two meristems are developmentally distinct, but few comparisons are available for other species.

In other cases, whole-genome sequences show that critical proteins that are critical for one species are simply absent in others. For example, RA1 and RA3 are not present in genomes of species of the BOP clade, implying that their function in rapid transition to a terminal spikelet in the short-branch (spikelet pair) meristems of maize and sorghum may be species- or clade specific. Genes that are genetically downstream of HvRA2 also differ from those genetically downstream of RA2 in maize, suggesting that each gene network may be only applicable in close relatives of barley or maize, respectively. Such presence-absence variation is only beginning to be explored.

In the future, we can anticipate identifying additional regulatory networks that make grass inflorescences so similar as well as the network components that make individual species morphologically distinct. The conserved components may be expected in all grasses, including orphan crops, whereas the variable components await analysis in disparate species.

Supplemental data

The following material is available in the online version of this article.

Supplemental Table S1. Protein-coding genes discussed in this article.

Acknowledgments

I thank the editors for the invitation to submit this review to the Mendel anniversary issue of the journal. Clint Whipple,

Madelaine Bartlett, and three anonymous reviewers provided valuable feedback on the ideas discussed here.

Funding

Work in the Kellogg lab has been supported by the National Science Foundation.

Conflict of interest statement. None declared.

References

- Alonso-Peral MM, Oliver SN, Casao MC, Greenup AA, Trevaskis B (2011) The promoter of the cereal *VERNALIZATION1* gene is sufficient for transcriptional induction by prolonged cold. *PLoS One* 6: e29456
- Arora R, Agarwal P, Ray S, Singh AK, Singh VP, Tyagi AK, Kapoor S (2007) MADS-box gene family in rice: genome-wide identification, organization and expression profiling during reproductive development and stress. *BMC Genom* 8: 242
- Bai S, Hong J, Li L, Su S, Li Z, Wang W, Zhang F, Liang W, Zhang D (2021) Dissection of the genetic basis of rice panicle architecture using a genome-wide association study. *Rice (NY)* 14: 77
- Bai X, Huang Y, Hu Y, Liu H, Zhang B, Smaczniak C, Hu G, Han Z, Xing Y (2017) Duplication of an upstream silencer of *FZP* increases grain yield in rice. *Nat Plants* 3: 885–893
- Bartlett ME, Thompson B (2014) Meristem identity and phyllotaxis in inflorescence development. *Front Plant Sci* 5: 508
- Bennetzen JL, Freeling M (1993) Grasses as a single genetic system: genome composition, collinearity and complementarity. *Trend Genet* 9: 259–261
- Bombliès K, Wang RL, Ambrose BA, Schmidt RJ, Meeley RB, Doebley J (2003) Duplicate *FLORICAULA/LEAFY* homologs *zfl1* and *zfl2* control inflorescence architecture and flower patterning in maize. *Development* 130: 2385–2395
- Bommert P, Satoh-Nagasawa N, Jackson D, Hirano HY (2005) Genetics and evolution of grass inflorescence and flower development. *Plant Cell Physiol* 46: 69–78
- Bommert P, Whipple C (2018) Grass inflorescence architecture and meristem determinacy. *Seminars Cell Devel Biol* 79: 37–47
- Bortiri E, Chuck G, Vollbrecht E, Rocheford T, Martienssen R, Hake S (2006) *ramosa2* encodes a LATERAL ORGAN BOUNDARY domain protein that determines the fate of stem cells in branch meristems of maize. *Plant Cell* 18: 574–585
- Butzin R (1979) Apikale reduktionen im infloreszenzbereich der gramineae. *Willdenowia* 9: 161–167
- Chae E, Tan QK, Hill TA, Irish VF (2008) An Arabidopsis F-box protein acts as a transcriptional co-factor to regulate floral development. *Development* 135: 1235–1245
- Chandler JW; (2018) Class VIIIb APETALA2 ethylene response factors in plant development. *Trends Plant Sci* 23: 151–162
- Chuck G, Cigan AM, Saeteurn K, Hake S (2007) The heterochronic maize mutant *Corngrass1* results from overexpression of a tandem microRNA. *Nat Genet* 39: 544–549
- Chuck G, Meeley R, Hake S (2008) Floral meristem initiation and meristem cell fate are regulated by the maize AP2 genes *ids1* and *sid1*. *Development* 135: 3013–3019
- Chuck G, Meeley R, Irish E, Sakai H, Hake S (2007) The maize *tas-elseed4* microRNA controls sex determination and meristem cell fate by targeting *Tasseled6/indeterminate spikelet1*. *Nat Genet* 39: 1517–1521
- Chuck G, Muszynski M, Kellogg E, Hake S, Schmidt RJ (2002) The control of spikelet meristem identity by the *branched silkless1* gene in maize. *Science* 298: 1238–1241
- Chuck GS, Brown PJ, Meeley R, Hake S (2014) Maize SBP-box transcription factors *unbranched2* and *unbranched3* affect yield traits by regulating the rate of lateral primordia initiation. *Proc Natl Acad Sci USA* 111: 18775–18780
- Claeys H, Vi SL, Xu X, Satoh-Nagasawa N, Eveland AL, Goldshmidt A, Feil R, Beggs GA, Sakai H, Brennan RG, et al. (2019) Control of meristem determinacy by trehalose 6-phosphate phosphatases is uncoupled from enzymatic activity. *Nat Plants* 5: 352–357
- Debernardi JM, Lin H, Chuck G, Faris JD, Dubcovsky J (2017) *microRNA172* plays a crucial role in wheat spike morphogenesis and grain threshability. *Development* 144: 1966–1975
- Demesa-Arevalo E, Abraham-Juarez MJ, Xu X, Bartlett M, Jackson D (2021) Maize RAMOSA3 accumulates in nuclear condensates enriched in RNA POLYMERASE II isoforms during the establishment of axillary meristem determinacy. *bioRxiv* 10.1101/2021.04.06.438639v1.full.pdf (April 6, 2021)
- Derbyshire P, Byrne ME (2013) MORE SPIKELETS1 is required for spikelet fate in the inflorescence of *Brachypodium*. *Plant Physiol* 161: 1291–1302
- Distelfeld A, Li C, Dubcovsky J (2009) Regulation of flowering in temperate cereals. *Curr Opin Plant Biol* 12: 178–184
- Dobrovolskaya O, Pont C, Sibout R, Martinek P, Badaeva E, Murat F, Chossion A, Watanabe N, Prat E, Gautier N, et al. (2015) FRIZZY PANICLE drives supernumerary spikelets in bread wheat. *Plant Physiol* 167: 189–199
- Doust A (2007) Architectural evolution and its implications for domestication in grasses. *Ann Bot* 100: 941–950
- Doust AN, Kellogg EA (2002) Inflorescence diversification in the panicoid "bristle grass" clade (Paniceae, Poaceae): evidence from molecular phylogenies and developmental morphology. *Am J Bot* 89: 1203–1222
- Doust AN, Mauro-Herrera M, Hodge JG, Stromski J (2017) The *C₄* model grass *Setaria* is a short day plant with secondary long day genetic regulation. *Front Plant Sci* 8: 1062
- Eveland AL, Jackson DP (2012) Sugars, signalling, and plant development. *J Exp Bot* 63: 3367–3377
- FAO (Food and Agriculture Organization of the United Nations) (2003) World Agriculture: Towards 2015–2030. Earthscan Publications Ltd, London
- Fujishiro Y, Agata A, Ota S, Ishihara R, Takeda Y, Kunishima T, Ikeda M, Kyoizuka J, Hobo T, Kitano H (2018) Comprehensive panicle phenotyping reveals that *qSrn7/FZP* influences higher-order branching. *Sci Rep* 8: 12511
- Gallavotti A, Long JA, Stanfield S, Yang X, Jackson D, Vollbrecht E, Schmidt RJ (2010) The control of axillary meristem fate in the maize *ramosa* pathway. *Development* 137: 2849–2856
- Gallavotti A, Malcomber S, Gaines C, Stanfield S, Whipple C, Kellogg E, Schmidt RJ (2011) BARREN STALK FASTIGIATE1 is an AT-hook protein required for the formation of maize ears. *Plant Cell* 23: 1756–1771
- Gallavotti A, Yang Y, Schmidt RJ, Jackson D (2008) The relationship between auxin transport and maize branching. *Plant Physiol* 147: 1913–1923
- Gallavotti A, Zhao Q, Kyoizuka J, Meeley RB, Ritter MK, Doebley JF, Pe ME, Schmidt RJ (2004) The role of *barren stalk1* in the architecture of maize. *Nature* 432: 630–635
- Galli M, Liu Q, Moss BL, Malcomber S, Li W, Gaines C, Federici S, Roshkovan J, Meeley R, Nemhauser JL, et al. (2015) Auxin signaling modules regulate maize inflorescence architecture. *Proc Natl Acad Sci USA* 112: 13372–13377
- Gao X, Liang W, Yin C, Ji S, Wang H, Su X, Guo C, Kong H, Xue H, Zhang D (2010) The *SEPALLATA*-like gene *OsMADS34* is required for rice inflorescence and spikelet development. *Plant Physiol* 153: 728–740
- Gao XQ, Wang N, Wang XL, Zhang XS (2019) Architecture of wheat inflorescence: insights from rice. *Trends Plant Sci* 24: 802–809
- Gauley A, Boden SA (2019) Genetic pathways controlling inflorescence architecture and development in wheat and barley. *J Integr Plant Biol* 61: 296–309

- Gocal GF, King RW, Blundell CA, Schwartz OM, Andersen CH, Weigel D (2001) Evolution of floral meristem identity genes. Analysis of *Lolium temulentum* genes related to *APETALA1* and *LEAFY* of Arabidopsis. *Plant Physiol* **125**: 1788–1801
- GPWG (Grass Phylogeny Working Group) (2001) Phylogeny and subfamilial classification of the Poaceae. *Ann Missouri Bot Garden* **88**: 373–457
- GPWG II (Grass Phylogeny Working Group II) (2012) New grass phylogeny resolves deep evolutionary relationships and discovers *C₄* origins. *New Phytologist* **193**: 304–312.
- Griffith DM, Osborne CP, Edwards EJ, Bachle S, Beerling DJ, Bond WJ, Gallaher TJ, Helliher BR, Lehmann CER, Leatherman L, et al. (2020) Lineage-based functional types: characterising functional diversity to enhance the representation of ecological behaviour in land surface models. *New Phytol* **228**: 15–23
- Harder LD, Prusinkiewicz P (2013) The interplay between inflorescence development and function as the crucible of architectural diversity. *Ann Bot* **112**: 1477–1493
- Harrop TWR, Mantegazza O, Luong AM, Bethune K, Lorieux M, Jouannic S, Adam H (2019) A set of *AP2*-like genes is associated with inflorescence branching and architecture in domesticated rice. *J Exp Bot* **70**: 5617–5629
- Houston K, Druka A, Bonar N, Macaulay M, Lundqvist U, Franckowiak J, Morgante M, Stein N, Waugh R (2012) Analysis of the barley bract suppression gene *Trd1*. *Theor Appl Genet* **125**: 33–45
- Hu L, Chen W, Yang W, Li X, Zhang C, Zhang X, Zheng L, Zhu X, Yin J, Qin P, et al. (2021) *OsSPL9* regulates grain number and grain yield in rice. *Front Plant Sci* **12**: 682018
- Huang P, Jiang H, Zhu C, Barry K, Jenkins J, Sandor L, Schmutz J, Box MS, Kellogg EA, Brutnell TP (2017) *Sparse panicle1* is required for inflorescence development in *Setaria viridis* and maize. *Nat Plants* **3**: 17054
- Huang X, Yang S, Gong J, Zhao Q, Feng Q, Zhan Q, Zhao Y, Li W, Cheng B, Xia J, et al. (2016) Genomic architecture of heterosis for yield traits in rice. *Nature* **537**: 629–633
- Huang Y, Zhao S, Fu Y, Sun H, Ma X, Tan L, Liu F, Sun X, Sun H, Gu P, et al. (2018) Variation in the regulatory region of *FZP* causes increases in secondary inflorescence branching and grain yield in rice domestication. *Plant J* **96**: 716–733
- Ikeda K, Ito M, Nagasawa N, Kyoizuka J, Nagato Y (2007) Rice *ABERRANT PANICLE ORGANIZATION1*, encoding an F-box protein, regulates meristem fate. *Plant J* **51**: 1030–1040
- Ikeda K, Nagasawa N, Nagato Y (2005) *ABERRANT PANICLE ORGANIZATION1* temporally regulates meristem identity in rice. *Dev Biol* **282**: 349–360
- Ikeda-Kawakatsu K, Maekawa M, Izawa T, Itoh J, Nagato Y (2012) *ABERRANT PANICLE ORGANIZATION 2/RFL*, the rice ortholog of Arabidopsis *LEAFY*, suppresses the transition from inflorescence meristem to floral meristem through interaction with *APO1*. *Plant J* **69**: 168–180
- Ikeda-Kawakatsu K, Yasuno N, Oikawa T, Iida S, Nagato Y, Maekawa M, Kyoizuka J (2009) Expression level of *ABERRANT PANICLE ORGANIZATION1* determines rice inflorescence form through control of cell proliferation in the meristem. *Plant Physiol* **150**: 736–747
- Jiao Y, Wang Y, Xue D, Wang J, Yan M, Liu G, Dong G, Zeng D, Lu Z, Zhu X, et al. (2010) Regulation of *OsSPL14* by *OsmiR156* defines ideal plant architecture in rice. *Nat Genet* **42**: 541–544
- Jin Y, Luo Q, Tong H, Wang A, Cheng Z, Tang J, Li D, Zhao X, Li X, Wan J, et al. (2011) An AT-hook gene is required for palea formation and floral organ number control in rice. *Dev Biol* **359**: 277–288
- Judziewicz EJ, Clark LG, Londoño X, Stern MJ (1999) American Bamboos. Smithsonian Institution Press, Washington, DC
- Judziewicz EJ, Soderstrom TR (1989) Morphological, anatomical, and taxonomic studies in *Anomochloa* and *Streptochoeta* (Poaceae: Bambusoideae). *Smithson Contrib Bot* **68**: 1–52
- Karim MR, Hirota A, Kwiatkowska D, Tasaka M, Aida M (2009) A role for Arabidopsis *PUCHI* in floral meristem identity and bract suppression. *Plant Cell* **21**: 1360–1372
- Kellogg EA (2000) A model of inflorescence development. In KL Wilson, DA Morrison, eds, *Monocots: Systematics and Evolution*. CSIRO, Melbourne, Australia, pp 84–88
- Kellogg EA (2007) Floral displays: genetic control of grass inflorescences. *Curr Opin Plant Biol* **10**: 26–31
- Kellogg EA (2015) Poaceae. In K Kubitzki, ed, *The Families and Genera of Vascular Plants*. Springer, Berlin, Germany, pp 1–416
- Kellogg EA, Camara PEAS, Rudall PJ, Ladd P, Malcomber ST, Whipple CJ, Doust AN (2013) Early inflorescence development in the grasses (Poaceae). *Front Plant Sci* **4**: 250
- Kim S, Soltis PS, Wall K, Soltis DE (2006) Phylogeny and domain evolution in the *APETALA2*-like gene family. *Mol Biol Evol* **23**: 107–120
- Kobayashi K, Maekawa M, Miyao A, Hirochika H, Kyoizuka J (2010) *PANICLE PHYTOMER2 (PAP2)*, encoding a *SEPALLATA* subfamily MADS-box protein, positively controls spikelet meristem identity in rice. *Plant Cell Physiol* **51**: 47–57
- Kobayashi K, Yasuno N, Sato Y, Yoda M, Yamazaki R, Kimizu M, Yoshida H, Nagamura Y, Kyoizuka J (2012) Inflorescence meristem identity in rice is specified by overlapping functions of three *API1/FUL*-like MADS box genes and *PAP2*, a *SEPALLATA* MADS box gene. *Plant Cell* **24**: 1848–1859
- Komatsu K, Maekawa M, Ujiie S, Satake Y, Furutani I, Okamoto H, Shimamoto K, Kyoizuka J (2003a) *LAX* and *SPA*: major regulators of shoot branching in rice. *Proc Natl Acad Sci USA* **100**: 11765–11770
- Komatsu M, Chujo A, Nagato Y, Shimamoto K, Kyoizuka J (2003b) *FRIZZY PANICLE* is required to prevent the formation of axillary meristems and to establish floral meristem identity in rice spikelets. *Development* **130**: 3841–3850
- Komatsuda T, Pourkheirandish M, He C, Azhaguvel P, Kanamori H, Perovic D, Stein N, Graner A, Wicker T, Tagiri A, et al. (2007) Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class homeobox gene. *Proc Natl Acad Sci USA* **104**: 1424–1429
- Koppolu R, Anwar N, Sakuma S, Tagiri A, Lundqvist U, Pourkheirandish M, Rutten T, Seiler C, Himmelbach A, Ariyadasa R, et al. (2013) *Six-rowed spike4 (Vrs4)* controls spikelet determinacy and row-type in barley. *Proc Natl Acad Sci USA* **110**: 13198–13203
- Kuijter HNJ, Shirley NJ, Khor SF, Shi J, Schwerdt J, Zhang D, Li G, Burton RA (2021) Transcript profiling of *MIKCC* MADS-Box genes reveals conserved and novel roles in barley inflorescence development. *Front Plant Sci* **12**: 705286
- Kyoizuka J (2014) Grass inflorescence: basic structure and diversity. *Adv Bot Res* **72**: 191–219
- Lee DY, An G (2012) Two *AP2* family genes, *SUPERNUMERARY BRACT (SNB)* and *OsINDETERMINATE SPIKELET1 (OsIDS1)*, synergistically control inflorescence architecture and floral meristem establishment in rice. *Plant J* **69**: 445–461
- Lee JH, Yoo SJ, Park SH, Hwang I, Lee JS, Ahn JH (2007) Role of *SVP* in the control of flowering time by ambient temperature in Arabidopsis. *Genes Dev* **21**: 397–402
- Lee YS, An G (2015) Regulation of flowering time in rice. *J Plant Biol* **58**: 353–360.
- Lehmann CER, Griffith DM, Simpson KJ, Anderson TM, Archibald S, Beerling DJ, Bond WJ, Denton E, Edwards EJ, Forrester EJ, et al. (2019) Functional diversification enabled grassy biomes to fill global climate space. *bioRxiv* doi: <https://doi.org/10.1101/583625> (March 21, 2019)
- Li C, Lin H, Chen A, Lau M, Jernstedt J, Dubcovsky J (2019a) Wheat *VRN1*, *FUL2* and *FUL3* play critical and redundant roles in spikelet development and spike determinacy. *Development* **146**: dev175398

- Li G, Kuijter HNJ, Yang X, Liu H, Shen C, Shi J, Betts N, Tucker MR, Liang W, Waugh R, et al. (2021a) MADS1 maintains barley spike morphology at high ambient temperatures. *Nat Plants* **7**: 1093–1107
- Li G, Zhang H, Li J, Zhang Z, Li Z (2021b) Genetic control of panicle architecture in rice. *Crop J* **9**: 590–597
- Li K, Debernardi JM, Li C, Lin H, Zhang C, Jernstedt J, Korff MV, Zhong J, Dubcovsky J (2021c) Interactions between SQUAMOSA and SHORT VEGETATIVE PHASE MADS-box proteins regulate meristem transitions during wheat spike development. *Plant Cell* **33**: 3621–3644
- Li Q, Wang Y, Wang F, Guo Y, Duan X, Sun J, An H (2016) Functional conservation and diversification of *APETALA1/FRUITFULL* genes in *Brachypodium distachyon*. *Physiol Plant* **157**: 507–518
- Li XY, Qian Q, Fu ZM, Wang YH, Xiong GS, Zeng DL, Wang XQ, Liu XF, Teng S, Hiroshi F (2003) Control of tillering in rice. *Nature* **422**: 618–621
- Li Y, Zhu J, Wu L, Shao Y, Wu Y, Mao C (2019b) Functional divergence of PIN1 paralogous genes in rice. *Plant Cell Physiol* **60**: 2720–2732
- Litt A (2007) An evaluation of A-function: evidence from the *APETALA1* and *APETALA2* gene lineages. *Int J Plant Sci* **168**: 73–91
- Litt A, Irish VF (2003) Duplication and diversification in the *APETALA1/FRUITFULL* floral homeotic gene lineage: implications for the evolution of floral development. *Genetics* **165**: 821–833
- Liu JM, Mei Q, Xue CY, Wang ZY, Li DP, Zhang YX, Xuan YH (2021) Mutation of G-protein gamma subunit *DEP1* increases planting density and resistance to sheath blight disease in rice. *Plant Biotechnol J* **19**: 418–420
- Liu Q, Peterson PM, Columbus JT, Zhao N, Hao G, Zhang D (2007) Inflorescence diversification in the "finger millet clade" (Chloridoideae, Poaceae): a comparison of molecular phylogeny and developmental morphology. *Am J Bot* **94**: 1230–1247
- Lu H, Dai Z, Li L, Wang J, Miao X, Shi Z (2017) OsRAMOSA2 shapes panicle architecture through regulating pedicel length. *Front Plant Sci* **8**: 1538
- Ma PF, Liu YL, Jin GH, Liu JX, Wu H, He J, Guo ZH, Li DZ (2021) The *Pharus latifolius* genome bridges the gap of early grass evolution. *Plant Cell* **33**: 846–864
- Matthes MS, Best NB, Robil JM, Malcomber S, Gallavotti A, McSteen P (2019) Auxin EvoDevo: conservation and diversification of genes regulating auxin biosynthesis, transport, and signaling. *Mol Plant* **12**: 298–320
- McKim SM (2020) Moving on up - controlling internode growth. *New Phytol* **226**: 672–678
- McSteen P, Malcomber S, Skirpan A, Lunde C, Wu X, Kellogg E, Hake S (2007) *barren inflorescence2* encodes a co-ortholog of the PINOID serine/threonine kinase and is required for organogenesis during inflorescence and vegetative development in maize. *Plant Physiol* **144**: 1000–1011
- Miura K, Ikeda M, Matsubara A, Song XJ, Ito M, Asano K, Matsuoka M, Kitano H, Ashikari M (2010) OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nat Genet* **42**: 545–549
- Moncur MW (1981) *Floral Initiation in Field Crops: An Atlas of Scanning Electron Micrographs*. CSIRO, Melbourne, Australia
- Nakagawa M, Shimamoto K, Kyoizuka J (2002) Overexpression of *RCN1* and *RCN2*, rice *TERMINAL FLOWER/CENTRORADIALIS* homologs, confers delay of phase transition and altered panicle morphology in rice. *Plant J* **29**: 743–750
- O'Connor DL, Runions A, Sluis A, Bragg J, Vogel JP, Prusinkiewicz P, Hake S (2014) A division in PIN-mediated auxin patterning during organ initiation in grasses. *PLoS Comput Biol* **10**: e1003447
- Oikawa T, Kyoizuka J (2009) Two-step regulation of LAX PANICLE1 protein accumulation in axillary meristem formation in rice. *Plant Cell* **21**: 1095–1108
- Patil V, McDermott HI, McAllister T, Cummins M, Silva JC, Mollison E, Meikle R, Morris J, Hedley PE, Waugh R, et al. (2019) *APETALA2* control of barley internode elongation. *Development* **146**: dev170373
- Phillips KA, Skirpan AL, Liu X, Christensen A, Slewinski TL, Hudson C, Barazesh S, Cohen JD, Malcomber S, McSteen P (2011) *vanishing tassels2* encodes a grass-specific tryptophan aminotransferase required for vegetative and reproductive development in maize. *Plant Cell* **23**: 550–566
- Pilatti V, Muchut SE, Uberti-Manassero NG, Vegetti AC, Reinheimer R (2019) Comparative study of the inflorescence, spikelet and flower development in species of Cynodonteae (Chloridoideae, Poaceae). *Bot J Linn Soc* **189**: 353–377
- Poethig RS (2009) Small RNAs and developmental timing in plants. *Curr Opin Genet Dev* **19**: 374–378
- Poursarebani N, Seidensticker T, Koppolu R, Trautewig C, Gawronski P, Bini F, Govind G, Rutten T, Sakuma S, Tagiri A, et al. (2015) The genetic basis of composite spike form in barley and 'miracle-wheat'. *Genetics* **201**: 155–165
- POWO (2021) Plants of the world online. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/> Retrieved 29 November 2021
- Preston JC, Christensen A, Malcomber ST, Kellogg EA (2009) MADS-box gene expression and implications for developmental origins of the grass spikelet. *Am J Bot* **96**: 1419–1429
- Preston JC, Kellogg EA (2006) Reconstructing the evolutionary history of paralogous *APETALA1/FRUITFULL*-like genes in grasses (Poaceae). *Genetics* **174**: 421–437
- Preston JC, Kellogg EA (2007) Conservation and divergence of *APETALA1/FRUITFULL*-like gene function in grasses: evidence from gene expression analyses. *Plant J* **52**: 69–81
- Preston JC, Kellogg EA (2008) Discrete developmental roles for temperate cereal grass *AP1/FUL*-like genes in flowering competency and the transition to flowering. *Plant Physiol* **146**: 265–276
- Preston JC, Wang H, Doebley J, Kellogg EA (2012) The role of *teosinte glume architecture (tga1)* in coordinated regulation and evolution of grass glumes and inflorescence axes. *New Phytol* **193**: 204–215
- Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E (2007) Evolution and development of inflorescence architectures. *Science* **316**: 1452–1456
- Qi J, Qian Q, Bu Q, Li S, Chen Q, Sun J, Liang W, Zhou Y, Chu C, Li X, et al. (2008) Mutation of the rice *Narrow leaf1* gene, which encodes a novel protein, affects vein patterning and polar auxin transport. *Plant Physiol* **147**: 1947–1959
- Ream TS, Woods DP, Schwartz CJ, Sanabria CP, Mahoy JA, Walters EM, Kaeppler HF, Amasino RM (2014) Interaction of photoperiod and vernalization determines flowering time of *Brachypodium distachyon*. *Plant Physiol* **164**: 694–709
- Reinhardt D, Pesce ER, Stieger P, Mandel T, Baltensperger K, Bennett M, Traas J, Friml J, Kuhlmeier C (2003) Regulation of phyllotaxis by polar auxin transport. *Nature* **426**: 255–260
- Reinheimer R, Vegetti AC (2008) Inflorescence diversity and evolution in the PCK clade (Poaceae: Panicoideae: Paniceae). *Plant Syst Evol* **275**: 133–167
- Reinheimer R, Vegetti AC, Rua GH (2013) Macroevolution of panicoid inflorescences: a history of contingency and order of trait acquisition. *Ann Bot* **112**: 1613–1628
- Reinheimer R, Zuloaga FO, Vegetti AC, Pozner R (2009) Diversification of inflorescence development in the PCK clade (Poaceae: Panicoideae: Paniceae). *Am J Bot* **96**: 549–564
- Ren D, Hu J, Xu Q, Cui Y, Zhang Y, Zhou T, Rao Y, Xue D, Zeng D, Zhang G, et al. (2018) FZP determines grain size and sterile lemma fate in rice. *J Exp Bot* **69**: 4853–4866
- Richardson A, Hake S (2022) The power of classic maize mutants: driving forward our fundamental understanding of plants. *Plant Cell* **34**: 2518–2533

- Saarela JM, Burke SV, Wysocki WP, Barrett MD, Clark LG, Craine JM, Peterson PM, Soreng RJ, Vorontsova MS, Duvall MR (2018) A 250 plastome phylogeny of the grass family (Poaceae): topological support under different data partitions. *PeerJ* **6**: e4299
- Sajo MG, Pabón-Mora N, Jardim J, Stevenson DW, Rudall PJ (2012) Homologies of the flower and inflorescence in the early-divergent grass *Anomochloa* (Poaceae). *Am J Bot* **99**: 614–628
- Sakuma S, Golan G, Guo Z, Ogawa T, Tagiri A, Sugimoto K, Bernhardt N, Brassac J, Mascher M, Hensel G, et al. (2019) Unleashing floret fertility in wheat through the mutation of a homeobox gene. *Proc Natl Acad Sci USA* **116**: 5182–5187
- Sakuma S, Schnurbusch T (2020) Of floral fortune: tinkering with the grain yield potential of cereal crops. *New Phytol* **225**: 1873–1882
- Satoh-Nagasawa N, Nagasawa N, Malcomber S, Sakai H, Jackson D (2006) A trehalose metabolic enzyme controls inflorescence architecture in maize. *Nature* **441**: 227–230
- Schrager-Lavelle A, Klein H, Fisher A, Bartlett M (2017) Grass flowers: an untapped resource. *J Syst Evol* **55**: 525–541
- Seetharam AS, Yu Y, Belanger S, Clark LG, Meyers BC, Kellogg EA, Hufford MB (2021) The *Streptochaeta* genome and the evolution of the grasses. *Front Plant Sci* **12**: 710383
- Selva C, Shirley NJ, Houston K, Whitford R, Baumann U, Li G, Tucker MR (2021) HvLEAFY controls the early stages of floral organ specification and inhibits the formation of multiple ovaries in barley. *Plant J* **108**: 509–527
- Shen C, Li G, Dreni L, Zhang D (2021) Molecular control of carpel development in the grass family. *Front Plant Sci* **12**: 635500
- Shitsukawa N, Takagishi A, Ikari C, Takumi S, Murai K (2006) WFL, a wheat FLORICAULA/LEAFY ortholog, is associated with spikelet formation as lateral branch of the inflorescence meristem. *Genes Genet Syst* **81**: 13–20
- Skirpan A, Wu X, McSteen P (2008) Genetic and physical interaction suggest that BARREN STALK1 is a target of BARREN INFLORESCENCE2 in maize inflorescence development. *Plant J* **55**: 787–797
- Smith RS, Guyomarc'h S, Mandel T, Reinhardt D, Kuhlemeier C, Prusinkiewicz P (2006) A plausible model of phyllotaxis. *Proc Natl Acad Sci USA* **103**: 1301–1306
- Soreng RJ, Peterson PM, Romaschenko K, Davidse G, Teisher JK, Clark LG, Barbera P, Gillespie LJ, Zuloaga FO (2017) A worldwide phylogenetic classification of the Poaceae (Gramineae) II: an update and a comparison of two 2015 classifications. *J Syst Evol* **55**: 259–290
- Tabuchi H, Zhang Y, Hattori S, Omae M, Shimizu-Sato S, Oikawa T, Qian Q, Nishimura M, Kitano H, Xie H, et al. (2011) LAX PANICLE2 of rice encodes a novel nuclear protein and regulates the formation of axillary meristems. *Plant Cell* **23**: 3276–3287
- Tanaka W, Pautler M, Jackson D, Hirano HY (2013) Grass meristems II: inflorescence architecture, flower development and meristem fate. *Plant Cell Physiol* **54**: 313–324
- Tsai AYL, Gazzarrini S (2014) Trehalose-6-phosphate and SnRK1 kinases in plant development and signaling: the emerging picture. *Front Plant Sci* **5**: 119
- Vegetti AC, Weberling F (1996) The structure of the paracladial zone in Poaceae. *Taxon* **45**: 453–460
- Vollbrecht E, Springer PS, Goh L, Buckler ES, Martienssen R (2005) Architecture of floral branch systems in maize and related grasses. *Nature* **436**: 1119–1126
- Walsh J, Freeling M (2002) The *liguleless2* gene of maize functions during the transition from the vegetative to the reproductive shoot apex. *Plant J* **19**: 489–495
- Wang L, Sun S, Jin J, Fu D, Yang X, Weng X, Xu C, Li X, Xiao J, Zhang Q (2015) Coordinated regulation of vegetative and reproductive branching in rice. *Proc Natl Acad Sci USA* **112**: 15504–15509
- Wang SS, Chung CL, Chen KY, Chen RK (2020) A novel variation in the *FRIZZLE PANICLE* (FZP) gene promoter improves grain number and yield in rice. *Genetics* **215**: 243–252
- Weberling F (1989) Morphology of Flowers and Inflorescences. Cambridge University Press, Cambridge
- Wei H, Zhao Y, Xie Y, Wang H (2018) Exploiting SPL genes to improve maize plant architecture tailored for high-density planting. *J Exp Bot* **69**: 4675–4688
- Whipple CJ (2017) Grass inflorescence architecture and evolution: the origin of novel signaling centers. *New Phytol* **216**: 367–372
- Whipple CJ, Hall DH, DeBlasio S, Taguchi-Shiobara F, Schmidt RJ, Jackson DP (2010) A conserved mechanism of bract suppression in the grass family. *Plant Cell* **22**: 565–578
- Whipple CJ, Kebrom TH, Weber AL, Yang F, Hall D, Meeley R, Schmidt R, Doebley J, Brutnell TP, Jackson DP (2011) *grassy tillers1* promotes apical dominance in maize and responds to shade signals in the grasses. *Proc Natl Acad Sci USA* **108**: E506–512
- Woods D, Dong Y, Bouche F, Bednarek R, Rowe M, Ream T, Amasino R (2019) A florigen paralog is required for short-day vernalization in a pooid grass. *Elife* **8**: e42153
- Woods DP, Hope CL, Malcomber ST (2011) Phylogenomic analyses of the BARREN STALK1/LAX PANICLE1 (BA1/LAX1) genes and evidence for their roles during axillary meristem development. *Mol Biol Evol* **28**: 2147–2159
- Woods DP, McKeown MA, Dong Y, Preston JC, Amasino RM (2016) Evolution of VRN2/Ghd7-like genes in vernalization-mediated repression of grass flowering. *Plant Physiol* **170**: 2124–2135
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS (2009) The sequential action of *miR156* and *miR172* regulates developmental timing in Arabidopsis. *Cell* **138**: 750–759
- Xiao Y, Guo J, Dong Z, Richardson A, Patterson E, Mangrum S, Bybee S, Bertolini E, Bartlett M, Chuck G, et al. (2021) Boundary domain genes were recruited to suppress bract growth and promote branching in maize. *bioRxiv*, doi.org/10.1101/2021.10.05.463134 (October 5, 2021)
- Xie K, Wu C, Xiong L (2006) Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and *microRNA156* in rice. *Plant Physiol* **142**: 280–293
- Xing Y, Zhang Q (2010) Genetic and molecular bases of rice yield. *Annu Rev Plant Biol* **61**: 421–442
- Xu M, Tang D, Cheng X, Zhang J, Tang Y, Tao Q, Shi W, You A, Gu M, Cheng Z, et al. (2019) OsPINOID regulates stigma and ovule initiation through maintenance of the floral meristem by auxin signaling. *Plant Physiol* **180**: 952–965
- Xu Q, Yu H, Xia S, Cui Y, Yu X, Liu H, Zeng D, Hu J, Zhang Q, Gao Z, et al. (2020) The C2H2 zinc-finger protein LACKING RUDIMENTARY GLUME 1 regulates spikelet development in rice. *Sci Bull* **65**: 753–764
- Yamaguchi A, Wu MF, Yang L, Wu G, Poethig RS, Wagner D (2009) The *microRNA*-regulated SBP-Box transcription factor SPL3 is a direct upstream activator of LEAFY, FRUITFULL, and APETALA1. *Dev Cell* **17**: 268–278
- Yan L, Loukolanov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J (2003) Positional cloning of the wheat vernalization gene *VRN1*. *Proc Natl Acad Sci USA* **100**: 6263–6268
- Yang D, He N, Zheng X, Zhen Y, Xie Z, Cheng C, Huang F (2020) Cloning of *long sterile lemma* (*lsl2*), a single recessive gene that regulates spike germination in rice (*Oryza sativa* L.). *BMC Plant Biol* **20**: 561
- Yang J, Bertolini E, Braud M, Preciado J, Chepote A, Jiang H, Eveland AL (2021) The SvFUL2 transcription factor is required for inflorescence determinacy and timely flowering in *Setaria viridis*. *Plant Physiol* **187**: 1202–1220
- Yang J, Yuan Z, Meng Q, Huang G, Perin C, Bureau C, Meunier AC, Ingouff M, Bennett MJ, Liang W, et al. (2017) Dynamic regulation of auxin response during rice development revealed by newly established hormone biosensor markers. *Front Plant Sci* **8**: 256

- Yang Z, Wang X, Gu S, Hu Z, Xu H, Xu C (2008) Comparative study of *SBP-box* gene family in Arabidopsis and rice. *Gene* **407**: 1–11
- Yao FQ, Li XH, Wang H, Song YN, Li ZQ, Li XG, Gao XQ, Zhang XS, Bie XM (2021) Down-expression of *TaPIN1s* increases the tiller number and grain yield in wheat. *BMC Plant Biol* **21**: 443
- Yao H, Skirpan A, Wardell B, Matthes MS, Best NB, McCubbin T, Durbak A, Smith T, Malcomber S, McSteen P (2019) The *barren stalk2* gene is required for axillary meristem development in maize. *Mol Plant* **12**: 374–389
- Yoshida A, Sasao M, Yasuno N, Takagi K, Daimon Y, Chen R, Yamazaki R, Tokunaga H, Kitaguchi Y, Sato Y, et al. (2013) TAWAWA1, a regulator of rice inflorescence architecture, functions through the suppression of meristem phase transition. *Proc Natl Acad Sci USA* **110**: 767–772
- Yoshida A, Suzaki T, Tanaka W, Hirano HY (2009) The homeotic gene *long sterile lemma* (*G1*) specifies sterile lemma identity in the rice spikelet. *Proc Natl Acad Sci USA* **106**: 20103–20108
- Yoshikawa T, Ito M, Sumikura T, Nakayama A, Nishimura T, Kitano H, Yamaguchi I, Koshiba T, Hibara K, Nagato Y, et al. (2014) The rice *FISH BONE* gene encodes a tryptophan aminotransferase, which affects pleiotropic auxin-related processes. *Plant J* **78**: 927–936
- Yuan H, Qin P, Hu L, Zhan S, Wang S, Gao P, Li J, Jin M, Xu Z, Gao Q, et al. (2019) OsSPL18 controls grain weight and grain number in rice. *J Genet Genomics* **46**: 41–51
- Zhu C, Box MS, Thiruppathi D, Hu H, Yu Y, Doust AN, McSteen P, Kellogg EA (2021a) Pleiotropic and non-redundant effects of an auxin importer in Setaria and maize. *bioRxiv*, <https://doi.org/10.1101/2021.10.14.464408> (October 14, 2021)
- Zhu QH, Hoque MS, Dennis ES, Upadhyaya NM (2003) Ds tagging of BRANCHED FLORETLESS 1 (BFL1) that mediates the transition from spikelet to floret meristem in rice (*Oryza sativa* L). *BMC Plant Biol* **3**: 6
- Zhu QH, Upadhyaya NM, Gubler F, Helliwell CA (2009) Over-expression of *miR172* causes loss of spikelet determinacy and floral organ abnormalities in rice (*Oryza sativa*). *BMC Plant Biol* **9**: 149
- Zhu W, Yang L, Wu D, Meng Q, Deng X, Huang G, Zhang J, Chen X, Ferrandiz C, Liang W, et al. (2021b) Rice *SEPALLATA* genes *OsMADS5* and *OsMADS34* cooperate to limit inflorescence branching by repressing the *TERMINAL FLOWER1*-like gene *RCN4*. *New Phytol* **233**: 1682–1700